



King's Research Portal

DOI:

[10.1016/j.jaci.2017.04.034](https://doi.org/10.1016/j.jaci.2017.04.034)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Jensen-Jarolim, E., Turner, M. C., & Karagiannis, S. N. (2017). AllergoOncology: IgE- and IgG4-mediated immune mechanisms linking allergy with cancer and their translational implications. *Journal of Allergy and Clinical Immunology*. <https://doi.org/10.1016/j.jaci.2017.04.034>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

AllergoOncology: IgE- and IgG₄-mediated immune mechanisms linking allergy with cancer and their translational implications



Erika Jensen-Jarolim, MD,^{a,b} Michelle C. Turner, PhD,^{c,d,e,f} and Sophia N. Karagiannis, PhD^{g,h} Vienna, Austria; Barcelona and Madrid, Spain; Ottawa, Ontario, Canada; and London, United Kingdom

Key words: Allergy, cancer, tumor, tolerance, IgE, effector cells, antibody-dependent cell-mediated cytotoxicity, AllergoOncology, clinical oncology, oncoimmunology

Since the detection of specific IgE in allergy, its potential role in cancer has been investigated and prompted the definition of the field of AllergoOncology.¹ Most recent developments are collected in a position paper by the European Academy of Allergy and Clinical Immunology.²

IgE AND CANCER

IgE is an antibody with properties distinct from those of other isotypes, specifically in terms of its affinity for its cognate Fc receptor FcεRI. FcεRI-IgE immune complex formation can activate potent effector cells normally associated with acute and chronic allergic responses. IgE effector cells, such as eosinophils, mast cells, and macrophages, are also known to infiltrate tumors; tumor-associated tissue eosinophilia (TATE) or tumor-associated macrophages (TAMs, which can constitute up to 50% of a tumor mass and can be alternatively activated, M2) are characteristics of tumor inflammation. While intratumoral or stromal mast cells have been correlated with tumor promotion, signs of mast cell degranulation, normally associated with IgE-immune complex formation, have been correlated with a more favorable prognosis. Mast cells are a prominent source of the proinflammatory cytokine TNF-α, which is known to promote antitumor immunity. Other released mediators turn on acute (histamine) or chronic (eg, slow-reacting substance of anaphylaxis and cytokines)

inflammation and promote amplification of innate effector mechanisms. These cells can harbor cytotoxic and phagocytic potential, which could be directed against tumors.

The potential efficacy of IgE antibodies engineered to recognize tumor antigens is exemplified *in vitro* by using cell-based assays, suggesting that IgE directed against tumor antigens engenders antibody-dependent cell-mediated cytotoxicity (ADCC) by human monocytes, whereas, through its interaction with FcγRs, IgG₁ of the same antigen specificity can instruct the same cells to trigger antibody-dependent cell-mediated phagocytosis (ADCP). IgE anti-cancer antibodies engage subsets of FcεRI-expressing effector cells to mediate ADCC against tumors without the inhibitory Fc receptor signals known to limit IgG effector functions in the tumor microenvironment. IgE is cross-linked by densely packed tumor antigens but not by soluble monovalent antigens, forming tumor-associated molecular patterns (TAMPs) on a cancer cell surface and therefore triggering effector cell activation at sites where antitumor immunity is needed (Fig 1). Therefore it is tempting to speculate that IgE antibodies directed against tumor antigens can propagate alternative or complementary antitumor functions to those of clinically available IgG mAbs specific for tumor antigens, such as epidermal growth factor receptor (EGFR) and human epidermal growth factor receptor 2 (HER2). Prompted by promising preclinical studies in numerous *in vivo* models of cancer and in nonhuman primates, the first clinical trial of an antitumor IgE antibody in patients with cancer is ongoing (NCT02546921).

From ^athe Institute of Pathophysiology and Allergy Research, Center of Pathophysiology, Infectiology and Immunology, Medical University Vienna; ^bThe Interuniversity Messerli Research Institute, University of Veterinary Medicine Vienna, Medical University Vienna and University Vienna; ^cthe Barcelona Institute for Global Health (ISGlobal), Barcelona; ^dUniversitat Pompeu Fabra (UPF), Barcelona; ^eCIBER Epidemiología y Salud Pública (CIBERESP), Madrid; ^fthe McLaughlin Centre for Population Health Risk Assessment, University of Ottawa; and ^gSt John's Institute of Dermatology, Division of Genetics & Molecular Medicine, Faculty of Life Sciences and Medicine, and ^hthe NIHR Biomedical Research Centre at Guy's and St Thomas' Hospitals and King's College London, Guy's Hospital, King's College London.

Supported by the European Academy for Allergy and Clinical Immunology, as an initiative of its AllergoOncology Task Force. E.J.-J. was supported by the Austrian Science Fund FWF (grant SFB F4606-B28). S.N.K. was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) based at Guy's and St Thomas' NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. S.N.K. was also supported by the Medical Research Council (MR/L023091/1), Cancer Research UK (C30122/A11527; C30122/A15774), the Academy of Medical Sciences, Breakthrough Breast Cancer (147), and the CR UK/NIHR in England/DoH for Scotland, Wales, and Northern Ireland Experimental Cancer Medicine Centre (C10355/A15587). ISGlobal is a member of the CERCA Programme, Generalitat de Catalunya.

Disclosure of potential conflict of interest: E. Jensen-Jarolim has received grants from the Austrian Science Fund (SFB F4606-B28), has received travel support from the

European Academy of Allergy and Clinical Immunology, has consultant arrangements with Allergy Therapeutics, and has received payment for lectures from Allergy Therapeutics and Thermo Fisher. M. C. Turner has consultant arrangements with ICF Incorporated. S. N. Karagiannis has received grants from the NIHR Biomedical Research Centre at Guy's and St Thomas's Hospitals NHS Trust and King's College London, the Medical Research Council, Cancer Research UK, the Academy of Medical Sciences, Breast Cancer Now, and CR UK/NIHR in England/DOH for Scotland, Wales, and Northern Ireland Experimental Cancer Medicine Centre and has received travel support from the European Academy of Allergy and Clinical Immunology.

Received for publication January 31, 2017; revised March 8, 2017; accepted for publication April 5, 2017.

Available online May 17, 2017.

Corresponding author: Erika Jensen-Jarolim, MD, Institute of Pathophysiology and Allergy Research, Center of Pathophysiology, Infectiology and Immunology, Medical University Vienna, Währinger G. 18-20, 1090 Vienna, Austria. E-mail: erika.jensen-jarolim@meduniwien.ac.at.

J Allergy Clin Immunol 2017;140:982-4.
0091-6749

© 2017 The Authors. Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.jaci.2017.04.034>




FIG 1. Antibody isotypes direct effector functions of innate immune cells infiltrating tumors. *Left*, Intratumoral IgG₄ supported by IL-10 and vascular endothelial growth factor (VEGF) produced by cancer and immune cells. IgG₄ occurs as intact IgG₄ antibody (a); can undergo Fab arm exchange (FAE), resulting in bispecific or half antibodies (b); or can interfere with IgG₁ Fc-mediated antitumor functions (c). IgG₄ might be unable to cross-link FcγRs on effector cells and engender antitumor functions. *Right*, Tumor antigen-specific IgE is shuttled through FcεRI⁺ effector cells into tumors. Overexpressed tumor antigens packed in lipid rafts assemble tumor-associated molecular patterns (TAMPs), leading to effective cross-linking of effector cell-bound IgE, followed by antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) of malignant cells (d); soluble monovalent tumor antigens do not lead to effector cell activation (e). Ag, Antigen; CDC, complement dependent cytotoxicity.

FROM T_H2 IMMUNE RESPONSES IN PATIENTS WITH ALLERGY AND ATOPY TO CANCER EPIDEMIOLOGY

Allergy is characterized by T_H2-dominant immune responses, featuring IL-4, IL-13, and thymic stromal lymphopoietin upregulation. Because these mediators can be observed in cancer tissues, strategies to activate these responses could in principle promote isotype switching to antitumor IgE and also IgE/FcεRI-associated cross-presentation by dendritic cells, resulting in activation of CD4⁺ T cells but also of CD8⁺ cytotoxic T lymphocytes. Therefore although T_H1 responses and a strong branch of CD8⁺ cytotoxic T lymphocytes are generally desired in oncology, activation of classical T_H2 cells has also been associated with improved survival. On the other hand, TATE and TAMs, especially those alternatively activated M2 populations, might be signs of a T_H2-biased immune response unable to restrict cancer growth. Evidence of an alternative T_H2 inflammatory milieu featuring enhanced IL-10 rather than IL-4 in patients with many cancers, including melanoma, can influence class-switching away from IgE and perhaps favor expression of

isotypes with more restricted effector functions, such as IgG₄ or IgA.

Therefore the question remains whether allergies in general can protect against cancer. Epidemiologic meta-analyses suggest a strong inverse association between allergy and atopy and risk of glioma, pancreatic cancer, and childhood leukemia, although there are limitations in previous studies related to measures of allergy history and latency period.³ Inverse associations of prediagnostic IgE and cancer risk have also been reported overall, as well as for glioma specifically,^{4,5} but fewer associations have been reported at other cancer sites.² Prospective studies in large cohorts are required to further understand the potential role of IgE and other immunologic parameters in cancer risk and potential underlying biological mechanisms to be able to better harness “classical” T_H2 immune responses and IgE against cancer.

IMMUNE TOLERANCE: GOOD OR BAD?

Immunotherapy strategies in oncology are hampered by an overwhelming immunosuppressive microenvironment in and

around tumors. Tumor cells and infiltrating immune cells, such as regulatory T cells and likely regulatory B cells,⁶ can secrete cytokines, such as TGF- β and IL-10, which support immune tolerance⁷ and moderate immune surveillance, hindering any antitumor activities of effector cells. Checkpoint inhibitors targeting programmed cell death protein 1 (PD-1) or cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) to unleash the cytotoxic potential of a patient's T-cell immunity are major breakthroughs in clinical oncology, even at the cost of severe systemic autoimmunity.

In contrast, allergen immunotherapy attempts to re-establish immune tolerance to the allergen. A hallmark of allergen immunotherapy is IgG₄, which is supported by IL-10-producing cells, such as regulatory T and regulatory B cells. IgG₄ antibodies can block allergic responses through various mechanisms, such as through interacting with the inhibitory Fc γ RIIb on B cells to block IgE synthesis.⁸ IgG₄ is also a special antibody isotype devoid of complement activation properties and with impaired effector functions. These characteristics might be due to hinge region sequences that render IgG₄ prone to Fab arm exchange and formation of naturally bispecific antibodies less able to be cross-linked by an allergen⁹ and therefore resistant to allergen-associated molecular patterns (AAMPs) (Fig 1).¹⁰ Furthermore, IgG₄ might repolarize M2a macrophages to the immunosuppressive phenotype M2b, which could be responsible for increased IL-10 secretion.¹¹ Strikingly, IgG₄ is expressed in tissues from patients with malignancies such as melanoma, in whom it can impair antitumor immunity and correlates with shorter survival and disease progression.¹² There is also increasing evidence to support positive correlations between IgG₄-related diseases, such as sclerosing cholangitis associated with autoimmune pancreatitis, with enhanced cancer risk,¹³ including more recent long-term follow-up investigations.¹⁴ However, the antigen specificities of IgG₄ antibodies are unclear.¹⁵ Notably, higher levels of IgE and IgG₄ recognizing the cancer antigens EGFR and HER2, but not carcinoembryonic antigen, were detected in the sera of patients with cancer compared with allergic patients.¹⁶ Therefore the tumor microenvironment could favor class-switching to IgG₄¹⁷ within a “modified T_H2 response” in a process that features high similarities to those reported in patients with cat allergy.¹⁸ Elucidating the conditions promoting IgG₄ isotype switching still requires the design of immunologically relevant animal models other than rodents, potentially monkeys or canines.²

Hence the opposite from allergen immunotherapy might be required in cancer treatment, ie, reduction of “alternative” T_H2 isotypes, such as IgG₄, *in situ* and simultaneous promotion of immune activatory IgE isotypes against cancer antigens. It might be envisaged that cancer immunotherapy strategies could aim at shifting existing IgG₄ responses to IgE through sequential isotype switching, perhaps with the aid of adjuvants.

CONCLUSION

Harnessing components of the “classical” T_H2 humoral immunity against tumors might offer novel oncological treatment avenues. Thus AllergoOncology might provide strategies that complement existing and emerging immuno-oncology therapies.

We thank Judit Fazekas for assistance with illustrations.

REFERENCES

- Jensen-Jarolim E, Achatz G, Turner MC, Karagiannis S, Legrand F, Capron M, et al. AllergoOncology: the role of IgE-mediated allergy in cancer. *Allergy* 2008;63:1255-66.
- Jensen-Jarolim E, Bax HJ, Bianchini R, Capron M, Corrigan C, Castells M, et al. AllergoOncology—the impact of allergy in oncology: EAACI position paper. *Allergy* 2017;72:866-87.
- Amirian ES, Zhou R, Wensch MR, Olson SH, Scheurer ME, Il'yasova D, et al. Approaching a scientific consensus on the association between allergies and glioma risk: a report from the Glioma International Case-Control Study. *Cancer Epidemiol Biomarkers Prev* 2016;25:282-90.
- Schwartzbaum J, Seweryn M, Holloman C, Harris R, Handelman SK, Rempala GA, et al. Association between prediagnostic allergy-related serum cytokines and glioma. *PLoS One* 2015;10:e0137503.
- Wulaningsih W, Holmberg L, Garmo H, Karagiannis SN, Ahlstedt S, Malmstrom H, et al. Investigating the association between allergen-specific immunoglobulin E, cancer risk and survival. *Oncoimmunology* 2016;5:e1154250.
- Mohr A, Renaudineau Y, Bagacean C, Pers JO, Jamin C, Bordron A. Regulatory B lymphocyte functions should be considered in chronic lymphocytic leukemia. *Oncoimmunology* 2016;5:e1132977.
- Stanic B, van de Veen W, Wirz OF, Ruckert B, Morita H, Sollner S, et al. IL-10-overexpressing B cells regulate innate and adaptive immune responses. *J Allergy Clin Immunol* 2015;135:771-80.e8.
- van de Veen W, Stanic B, Wirz OF, Jansen K, Globinska A, Akdis M. Role of regulatory B cells in immune tolerance to allergens and beyond. *J Allergy Clin Immunol* 2016;138:654-65.
- Davies AM, Sutton BJ. Human IgG₄: a structural perspective. *Immunol Rev* 2015;268:139-59.
- Pali-Scholl I, Jensen-Jarolim E. The concept of allergen-associated molecular patterns (AAMP). *Curr Opin Immunol* 2016;42:113-8.
- Jensen-Jarolim E, Roth-Walter F, Pacios L, Wagner S, Glenk L, Hofstetter G, et al. IgG₄ drives M2 macrophages to cortisol, Lcn-2 and IL-10 release: implications in maintenance of tolerance and allergen immunotherapy. Annual Meeting of the American Academy of Allergy, Asthma and Immunology (AAAAI). *J Allergy Clin Immunol* 2016;137(suppl):AB403, L43.
- Karagiannis P, Villanova F, Josephs DH, Correa I, Van Hemelrijck M, Hobbs C, et al. Elevated IgG₄ in patient circulation is associated with the risk of disease progression in melanoma. *Oncoimmunology* 2015;4:e1032492.
- Wallace ZS, Wallace CJ, Lu N, Choi HK, Stone JH. Association of IgG₄-related disease with history of malignancy. *Arthritis Rheumatol* 2016;68:2283-9.
- Asano J, Watanabe T, Oguchi T, Kanai K, Maruyama M, Ito T, et al. Association between immunoglobulin G₄-related disease and malignancy within 12 years after diagnosis: an analysis after longterm followup. *J Rheumatol* 2015;42:2135-42.
- Karagiannis P, Gilbert AE, Josephs DH, Ali N, Dodev T, Saul L, et al. IgG₄ subclass antibodies impair antitumor immunity in melanoma. *J Clin Invest* 2013;123:1457-74.
- Zennaro D, Capalbo C, Scala E, Liso M, Spillner E, Penichet M, et al. IgE, IgG₄ and IgG response to tissue-specific and environmental antigens in patients affected by cancer. Annual Meeting of the European Academy of Allergy and Clinical Immunology, Istanbul. *Allergy* 2011;66:100.
- Crescioli S, Correa I, Karagiannis P, Davies AM, Sutton BJ, Nestle FO, et al. IgG₄ Characteristics and Functions in Cancer Immunity. *Curr Allergy Asthma Rep* 2016;16:7.
- Renand A, Archila LD, McGinty J, Wambre E, Robinson D, Hales BJ, et al. Chronic cat allergen exposure induces a TH2 cell-dependent IgG₄ response related to low sensitization. *J Allergy Clin Immunol* 2015;136:1627-35, e1-13.